


# FROM VECTOR AND MATRICES TO TENSORS: COMMENTS & QUESTIONS

Carlos I. PÉREZ SÁNCHEZ, Univ. Heidelberg & ESI, Vienna  
 “Invitation to Recursion, Resurgence and Combinatorics”  
 OIST, Okinawa, Japan. April 2023

To rethink some of the geometric tools of ‘tensor models’ invites us:

“The construction of tensor models was motivated by the idea of generalizing to higher dimensions the familiar relation of matrix models to random two-dimensional geometries (...) The status of this program is unclear, since it is not clear that the rather special Feynman diagrams that are generated by (...)



describe a useful class of random (...) geometries” E. WITTEN *J. Phys. A* ‘16

Can we import (further) tools from matrix/vector models?

## 1. DISCRETE SURFACES

- To address the enumeration of surfaces constructed by ‘gluings of polygons’, we first address a simpler problem: count gluings of a rooted polygon of  $2p$  sides. By a gluing, we mean pairings  $\pi \in \mathcal{P}_2(2p)$  of its sides. We think of  $\pi$  as chords inside the polygon; ‘rooted’ means that the polygon is fixed while  $\mathbb{Z}_{2p}$  rotates the chord diagram
- from the  $(2p-1)!! = (2p)!/2^p p! = \#\mathcal{P}_2(2p)$  gluings, let  $c_g(p)$  be the number of those having genus  $g$ . Call  $Q_p(N)$  the generating series (a polynomial in this case) in the sense

$$Q_p(N) = \frac{1}{N^2} \sum_{g \geq 0} c_g(p) N^{2-2g},$$

where the scalings in  $N$  (still just a formal variable to be clarified) are by convenience. Notice that for  $g > 0$ ,  $c_g(p)$  are higher genus generalizations of the Catalan number  $c_0(p) = \frac{1}{p+1} \binom{2p}{p}$ . For instance,  $N^2 Q_3(N) = 5N^2 + 10N^0$

- a further step is dropping the restriction of the polygons having  $2p$ -sides and summing over the number of sides

$$1 + 2zN + 2z \sum_{p \geq 1} \frac{Q_p(N)}{(2p-1)!!} (Nz)^p = \left[ \frac{1+z}{1-z} \right]^N \quad (1)$$

J. HARER & D. ZAGIER *Invent. Math.* ‘86. This generating function contains all the information, since the coefficient  $[z^{p+1}, N^{p+1-2g}]_{\frac{1}{2}\text{RHS}}$  gives the genus- $g$  fraction of gluings of  $2p$ -agons for arbitrary  $p$

- a matrix integral representation was relevant in one of the many proofs of Eq. 1. With the trace  $\text{Tr}(H) = \sum_{a=1}^N H_{a,a}$ ,

$$Q_p(N) = \int_{M_N(\mathbb{C})_{\text{s.a.}}} \frac{1}{N} \text{Tr}(H^{2p}) d\mu(H) =: \langle \frac{1}{N} \text{Tr} H^{2p} \rangle_G,$$

where  $d\mu(H)$  is the normalized Gaussian measure  $d\mu(H) = K_N \exp[-(N/2) \text{Tr} H^2] dH$ . While in order to get Formula 1 one has to work more, the matrix integral representation is readily obtained via  $\langle H_{a,b} H_{c,d} \rangle_G = \frac{1}{N} \delta_{a,d} \delta_{b,c}$  and

$$\langle H_{a_1, b_1} \cdots H_{a_{2p}, b_{2p}} \rangle_G = \sum_{\pi \in \mathcal{P}_2(2p)} \prod_{(i,j) \in \pi} \langle H_{a_i, b_i} H_{a_j, b_j} \rangle_G$$

- if one allows connected ‘gluings’ of several polygons, the natural concept is *combinatorial map*  $G = (J, \phi, \tau)$ , where  $J = \{1, \dots, h\}$  is the set of  $h \in 2\mathbb{N}$  half-edges and  $\phi, \tau \in \mathfrak{S}_h = S_h$ , being  $\tau$  free from fixed points and  $\tau^2 = 1$ . The faces, edges and vertices of the map are the cycles (denoted  $\mathcal{C}$ ) of  $\phi, \tau$  and  $v = \phi \circ \tau$ , respectively. Thus  $\#\mathcal{C}(v) - \#\mathcal{C}(\tau) - \#\mathcal{C}(\phi) = \chi(G) = 2 - 2g$ . For instance,  $J = \{1, \dots, 6\}, \phi = (162435), \tau = (14)(25)(36)$  describe a map with  $\chi(\text{C}) = 0$ , since  $v = (132)(465)$
- to generate maps, one introduces a potential  $V(x) = \sum_{0 < k \leq d} t_k x^k / k$  which yields a new partition function  $\mathcal{Z} = C_N \int_{M_N(\mathbb{C})_{\text{s.a.}}} e^{-NV(H)} dH$ . Maps (with  $\partial$ ) are counted by

$$\langle \text{Tr} H^{\ell_1} \cdots \text{Tr} H^{\ell_n} \rangle =: \sum_{g \geq 0} N^{2-2g-n} \mathcal{T}_{\ell_1, \ell_2, \dots, \ell_n}^{(g)}$$

where the LHS is computed with  $\langle P(H) \rangle := \mathcal{Z}^{-1} \int P(H) e^{NV(H)} dH$ . These can be obtained when  $\partial_{\ell} := \partial / \partial t_{\ell}$  hits the partition function  $\mathcal{Z}$

- with  $\ell = (\ell_2, \dots, \ell_n)$ , *Tutte Equations*  $|_{t_1=0, t_2=-1}$  read

$$\mathcal{T}_{\ell_1+1, \ell}^{(g)} = \sum_{j=3}^d t_j \cdot \mathcal{T}_{\ell_1+j-1, \ell}^{(g)} + \sum_{c=2}^k \ell_c \cdot \mathcal{T}_{\ell_1+\ell_c-1, \ell_2, \dots, \widehat{\ell_c}, \dots, \ell_n}^{(g)} + \sum_{\substack{p, q \text{ with} \\ p+q=\ell_1-1}} \left\{ \sum_{\substack{h_1+h_2=g \\ I \cup J = \ell}} \mathcal{T}_{p, I}^{(h_1)} \times \mathcal{T}_{q, J}^{(h_2)} + \mathcal{T}_{p, q, \ell}^{(g-1)} \right\}$$

- to obtain these (with  $t_1, t_2 + 1 \neq 0$ ) one can use *Schwinger-Dyson equations* (SDE), sketched next: from  $\int d(X e^{-S/\hbar}) = 0$ , it holds  $\int [dX X - \frac{1}{\hbar} dS(X)] e^{-S/\hbar} = 0$ . So  $\langle \text{grad} S(X) \rangle = \hbar \langle \text{div} X \rangle$ . For  $\hbar \rightarrow 0$ , the SDE yield (classical EOM)

- Tutte Equations can be restated as differential operators,  $\mathcal{L}_k, k = -1, 0, 1, 2 \dots$  that annihilate the partition function  $\mathcal{Z}' = \exp(N^2 t_0) \mathcal{Z}, \mathcal{L}_k \mathcal{Z}' = 0$ . Omitting the cases  $k = 0, \pm 1, \mathcal{L}_k$  for  $k > 1$  is given by

$$\sum_{j=1}^{k-1} \frac{j(k-j)}{N^2} \partial_j \partial_{k-j} + \frac{2k}{N^2} \partial_k \partial_0 + \sum_{j \in \mathbb{N}} (j+k) t_j \partial_{j+k},$$

which satisfies the Witt algebra  $\mathfrak{w}, [\mathcal{L}_p, \mathcal{L}_q] = (p-q) \mathcal{L}_{p+q}$  (if  $\mathcal{L}_{-1, 0, 1}$  are added) for  $p, q \in \mathbb{Z}_{\geq -1}$  i.e. non-central vir

## 2. ‘GEOMETRY’ AND TENSOR MODELS

- Let’s begin by geometry: having gravitation as purpose, we are interested in PL-manifolds, as there the path integral seems better controlled. In fact, Alexander theorem—and refined (independent) versions by Ramirez, Montesinos and Hilden—states that connected, closed orientable 3-manifolds are *covers* over  $S^3$  branched along a link/knot (which is far from unique). This holds in general dimension  $D$ , the branching over  $S^D$  taking place always at codimension-2 subset of that sphere




Fig. 1 (L)  $\mathbb{T}^2$  as branched over  $S^2$  (C&R) Barycentric subdivision  $D = 2, 3$

- one can always ‘color’ those triangulations by barycentric subdivision. Tensor Models computes integrals of the form

$$\int B(\phi, \bar{\phi}) e^{-N^2 \bar{\phi}_{pqr} \phi_{pqr}} d\phi d\bar{\phi}$$

which are performed over functions  $\phi, \bar{\phi} : \{1, \dots, N\}^3 \rightarrow \mathbb{C}$  satisfying that for each argument  $1 \leq p, q, r \leq N$  the evaluation  $\phi_{pqr} = \phi(p, q, r)$  transforms independently under  $U(N)$ . Interesting for  $B(\phi, \bar{\phi})$  are  $U(N)^3$ -invariants, aka ‘bubbles’

$$\left( \begin{array}{c} 2 \\ \text{---} \\ 1 \\ \text{---} \\ 3 \end{array} \right) = \phi_{a_1 a_2 a_3} \bar{\phi}_{a_1 b_2 c_3} \phi_{b_1 b_2 b_3} \bar{\phi}_{b_1 a_2 b_3} \phi_{c_1 c_2 c_3} \bar{\phi}_{c_1 c_2 a_3}$$


which we rather specify via (regularly) 3-colored (vertex-bipartite) graphs

- those invariants form a basis for the interactions,  $S(\phi, \bar{\phi}) = \sum_{\mathcal{B}, U(N)^3\text{-inv.}} t_{\mathcal{B}} \mathcal{B}(\phi, \bar{\phi})$  being  $t_{\mathcal{B}}$  formal variables.

$$\mathcal{Z}_N = \int \exp[-N^2 S(\phi, \bar{\phi})] d\phi d\bar{\phi} \in \mathbb{C}[[\{t_{\mathcal{B}}\}_{\mathcal{B}, \text{tricolored}}]]$$

- ★ Q0: Is there a Harer-Zagier function for tensor integrals analogous to  $(1+z)^N/(1-z)^N$  for matrix integrals?

- amplitude of a Feynman graph  $\mathcal{G}$ , now having  $D+1$  colors, scales  $\sim N^{\#\text{faces}(\mathcal{G}) - D(D-1)\#\text{vertices}(\mathcal{G})/4}$  and interpreting  $\mathcal{G}$  as the gluing  $\Delta(\mathcal{G})$  of  $\#V(\mathcal{G})$  of equilateral  $D$ -simplices whose boundaries are glued following the edges of  $\mathcal{G}$




one can relate the amplitudes to Regge-Einstein-Hilbert action (which in some units reads)  $S_{\text{REH}}(\Delta(\mathcal{G})) = \sum_D \text{simplices } \sigma \text{ vol}(\sigma) - \sum_{\text{codim } 2 \text{ simpl. } \tau} \delta(\tau) \text{vol}(\tau)$ , being  $\delta(\tau)$  the deficiency angle at  $\tau$

- ★ Q1: While in  $D = 2$  the equilateral condition is irrelevant, in  $D \geq 3$  it seems only extremely convenient. Can one for  $D \geq 3$  find an association of geometric parameters  $P_D(a_1, \dots, a_D) \in \{\text{angles, areas, lengths, ...}\}$  that respects the  $1/N$  expansion?

- For invariants (bubbles)  $\mathcal{A}, \mathcal{B}$  and  $v \in V_{\bullet}(\mathcal{A}), w \in V_{\bullet}(\mathcal{B})$ ,  $\mathcal{A} * \mathcal{B} = \sum_{u \in V_{\bullet}(\mathcal{A})} (\mathcal{A} \sqcup \mathcal{B}) \setminus \{u, w\} \Big|_{\text{glue colorwise}}$  (rooted at  $v$ )

which in topologically boils down to connected sums:



- if  $\mathcal{A}_{v,w} = \mathcal{A} \setminus \{v, w\} \Big|_{\text{glued by color}}^{\text{broken edges}}$  for  $w \in V_{\circ}(\mathcal{A})$ , the loop or Schwinger-Dyson Equations can be expressed as  $\mathcal{L}_{\mathcal{A},v} \mathcal{Z}_N = 0$ , where


$$\mathcal{L}_{\mathcal{A},v} = \sum_{w \in V_{\circ}(\mathcal{A})} N^{\#\text{Edges}(w \leftrightarrow v)} \prod_{\rho \in \pi_0(\mathcal{A}_{v,w})} \left[ -\frac{1}{N^2} \frac{\partial}{\partial t_{\rho}} \right] + \sum_{\mathcal{B} \in \mathcal{S}} t_{\mathcal{B}} \frac{\partial}{\partial t_{\mathcal{B} * \mathcal{A}}}$$

so for connected  $\mathcal{A}$ ,  $\mathcal{L}_{\mathcal{A},v}$  can be of second degree, e.g. for  $\mathcal{A} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array}$ , if  $v, w$  are the middle vertices

- Similar graph operations describe the Polchinski equation T. KRAJEWSKI & R. TORIUMI *J. Phys. A* '16 (Wetterich's is w.i.p.)
- by R. GURÄU *Nucl. Phys. B* '12 these operators satisfy

$$[\mathcal{L}_{\mathcal{A},v}, \mathcal{L}_{\mathcal{B},w}] := \mathcal{L}_{\mathcal{A} * \mathcal{B},v} - \mathcal{L}_{\mathcal{B} * \mathcal{A},w}$$

- this Guräü algebra restricts to  $\mathfrak{w}_{\geq 1}$  algebra  $[\mathcal{L}_{C_p}, \mathcal{L}_{C_q}] = (p-q)\mathcal{L}_{C_{p+q}}$  for  $p, q \in \mathbb{Z}_{\geq 1}$  if we restrict to cycles,



- ★ Q2: Can one find a recursion (satisfied by the correlators) in tensor models, even though it is not topological?

### 3. AIRY STRUCTURES AND TOPOLOGICAL RECURSION (TR)

Airy structures M. KONTSEVICH & Y. SOIBELMAN '17 capture the essence of TR



- Let  $W^*$  be the vector space with basis  $\{t_j\}_{j=1}^d$ . If  $\hbar$  is a formal variable, a (quantum) Airy structure on  $W$  is a family of operators  $\{L_k\}_k$  on  $\text{Sym}(W^*)[[\hbar, \hbar^{-1}]]$  where  $L_k$  reads

$$-\frac{1}{2}(\mathbf{t}, A^k \mathbf{t}) - \hbar(\mathbf{t}, B^k \partial) - \frac{\hbar^2}{2}(\partial, C^k \partial) + \hbar(\partial_k - D_k)$$

and such that  $[L_i, L_j] = \hbar \sum_k f_{i,j}^k L_k$ , being  $f^k, A^k, B^k, C^k \in M_d(\mathbb{C})$ , where  $A^k$  and  $C^k$  are symmetric, while  $f^k$  is skew-symmetric for each  $k$  (not a matrix index nor exponent, abusing on notation)

- The Lie algebra condition implies that  $A$ , seen as a tensor, is fully symmetric; that  $f_{i,j}^k = B_{j,k}^i - B_{i,k}^j$ ; and three IHX-relations described next. To the six vertices one associates letters. Red edges have indices that run. Further, the indices of each letter  $O$  at the vertices  $O_{\square}^{\square}$  is determined in the sense of the arrow, starting at the shaded edge. The IHX-relation for  $(v_1, v_2, \dots, v_6) = (B, B, B, B, C, A)$  is


that  $\sum_{a=1}^d B_{j,a}^i A_{k,l}^a + B_{k,a}^i A_{a,l}^j + B_{l,a}^i A_{a,k}^j$  is  $(i \leftrightarrow j)$ -symmetric. Similar relations hold for  $(v_1, v_2, \dots, v_6) = (B, A, B, B, B, A)$  and  $(C, B, B, C, C, B)$ .



- THM (Kontsevich-Soibelman) There exists a unique  $\hbar^{-1}F \in \text{Sym}(W^*)[[\hbar]]$  such that  $\{L_j e^F = 0\}_{j=1, \dots, d}$ . Proof sketch of uniqueness (cf. G. BOROT *Rev. Math. Phys.* '20). Expand  $F = \sum_{g \geq 0} \hbar^{g-1} \sum_{n \geq 1} \sum_{I, \#I=n} F_{g,n}[I] t_I / n!$ , with  $I = (i_1, \dots, i_n) \in \{1, \dots, d\}^n$  and  $t_I = t_{i_1} \dots t_{i_n}$  in multi-index notation, and read off the coefficient of  $\hbar^g \times t_{i_2} \dots t_{i_n} / (n-1)!$  in  $\exp(-F) L_{i_1} \exp(F) = 0$ . This yields  $F_{0,3}[i_1, i_2, i_3] = A_{i_2, i_3}^{i_1}$  and  $F_{1,1}[i_1] = D_{i_1}$  for  $\chi = -1$ , while for higher  $-\chi$ ,  $F_{g,n}[i_1, i_2, \dots, i_n]$  is determined by recursion and equals (cf. last page)

$$\sum_{m=2}^n B_{i_m, a}^{i_1} F_{g, n-1}[a, i_2, \dots, \widehat{i_m}, \dots, i_n] \quad (i_q := \#J_q, q = 1, 2) + \frac{1}{2} C_{a,b}^{i_1} \left\{ F_{g-1, n+1}[a, b, i_2, \dots, i_n] \right. \quad (3)$$

$$\left. + \sum_{J_1 \cup J_2 = \{i_2, \dots, i_n\}}^{h_1 + h_2 = g} F_{h_1, 1+j_1}[a, J_1] \times F_{h_2, 1+j_2}[b, J_2] \right\}$$



- the boundaries above are not oriented; but, parenthetically, the ABCD-terms could stem from a TQFT  $\mathcal{F} : \text{Bord}_2 \rightarrow \text{Vect}_{\mathbb{C}}$   
 $A = \mathcal{F} \left( \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right), B = \mathcal{F} \left( \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right), C = \mathcal{F} \left( \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right), D = \mathcal{F} \left( \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right)$

4. THE VOLUME OF THE MODULI SPACE  $\mathcal{M}_{g,n}(L)$ 

- $\mathcal{T}_{g,n}(L) = \{\text{metrics on } \Sigma_{g,n} : \text{length of boundary } b_j = L_j\} / \{\text{conformal maps}\}$ , with  $L = (L_1, \dots, L_n)$
- $\Gamma_{g,n} = \{\text{Diff}(\Sigma_{g,n}) \text{ that keep labels}\} / \{\text{isotopies to id}_{\Sigma_{g,n}}\}$
- $\mathcal{M}_{g,n}(L) = \mathcal{T}_{g,n}(L) / \Gamma_{g,n} = \text{Teichmüller/mapping class}$
- decomposition of a stable surface  $\Sigma_{g,n}$  in simple closed curves yields  $p$   $Y$ -pieces, each having Euler number  $-1$ , so  $p = -\chi(\Sigma_{g,n})$ . From the  $3p$  geodesic boundaries,  $n$  are *not* glued, so there are  $\frac{1}{2}(p - n) = 3g + n - 3 := d_{g,n}$  inner pairings of cycles, whose lengths  $\ell_j$  can coincide. The twisting angle  $\theta_j$  of one cycle with respect to the other is another parameter
- $\{\ell_j, \theta_j\}_{j=1, \dots, 3g+n-3}$  are in fact the *Fenchel-Nielsen* coordinates of  $\mathcal{T}_{g,n}$ . The form  $\omega_{\text{wp}} = \sum_j d\ell_j \wedge d\theta_j$  is  $\Gamma_{g,n}$ -invariant, as shown by WOLPERT '85, and  $\omega_{\text{wp}}^{\wedge d_{g,n}} / d_{g,n}!$  defines the volume form of  $\mathcal{M}_{g,n}(L)$  and  $V_{g,n}(L) = \text{vol}[\mathcal{M}_{g,n}(L)]$

- MIRZAKHANI *JAMS* '07 TR states that  $V_{g,n+1}(L_0, L)$  equals

$$\sum_{m>0} \int_{\mathbb{R}_+} B_{\text{Mirz}}(L_0, L_m, \ell) V_{g,n}(\ell, L_1, \dots, \widehat{L_m}, \dots, L_n) d\ell \\ + \frac{1}{2} \int_{\mathbb{R}_+^2} C_{\text{Mirz}}(L_0, \ell, \ell') [V_{g-1, n+2}(\ell, \ell', L_1, \dots, L_n) \\ + \sum_{\substack{J_1 \cup J_2 = \{1, \dots, n\} \\ h_1 + h_2 = g}} V_{h_1, 1+j_1}(\ell, J_1) V_{h_2, 1+j_2}(\ell', J_2)] d\ell d\ell',$$

where  $B_{\text{Mirz}}(L_1, L_2, L_3)$  and  $C_{\text{Mirz}}(L_1, L_2, L_3)$  are given by

$$\frac{L_3}{L_1} \log \frac{[1 + e^{(L_3+L_2-L_1)/2}][1 + e^{(L_3-L_2-L_1)/2}]}{[1 + e^{(L_3+L_2+L_1)/2}][1 + e^{(L_3-L_2+L_1)/2}]}$$

$$\text{and } 2 \frac{L_2 L_3}{L_1} \log \frac{1 + e^{(L_3+L_2-L_1)/2}}{1 + e^{(L_3+L_2+L_1)/2}}, \text{ respectively}$$

- keeping the coefficients of the volumes as amplitudes,

$$V_{g,n}(L) = \sum_{a_1, \dots, a_n \geq 0} F_{g,n}[a_1, \dots, a_n] \prod_{j=1}^n L_j^{2a_j},$$

Mirzakhani's TR and (3) lead to the Airy structure:

$$B_{j,k}^i = \frac{(2k+1)!}{(2i+1)!(2j+1)!} (2j+1) \theta_{k-j-i} \quad (4a)$$

$$C_{j,k}^i = \frac{(2j+1)!(2k+1)!}{(2i+1)!} \theta_{k+j+1-i}, \quad (4b)$$

where  $\sum_{k+1 \geq 0} z^{2k} \theta_k / dz = 4\pi / \sin(2\pi z) dz^2 =: 1/y(z) dz^2$ . But we need the initial  $A$  and  $D$  terms

- the remarkable formula in M. KONTSEVICH *Anal. and Appl.*, '91 uses maps, or ribbon graphs, to compute also intersection numbers

$$\sum_{\substack{a_j \in \mathbb{Z}_{\geq 0}, \text{ for all } j \\ a_1 + \dots + a_n = \dim_{\mathbb{C}} \overline{\mathcal{M}}_{g,n}}} \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{a_1} \cdot \psi_2^{a_2} \cdots \psi_n^{a_n} \prod_{j=1}^n \frac{(2a_j - 1)!!}{\lambda_j^{2a_j+1}} \\ = \sum_{G \text{ trivalent, of topology } (g,n)} \frac{2^{2g-2+n}}{\#\text{Aut}(G)} \prod_{e \in E(G)} \frac{1}{\lambda_{L(e)} + \lambda_{R(e)}}$$

- e.g. there is a unique  $(1, 1)$ -graph:

$$G = \text{graph} \Rightarrow \frac{1}{\lambda^3} \int_{\overline{\mathcal{M}}_{1,1}} \psi_1 = \frac{2^1}{\#\text{Aut}(G)} \frac{1}{(2\lambda)^3}$$

$$\text{Aut}(G) = \{\psi \in \mathfrak{S}_6 : \text{commuting with } \phi \text{ and } \tau\}$$

$$= \{\text{id}, (123)(456), (132)(465), \tau, \phi, \phi^{-1}\}$$

then  $\int_{\overline{\mathcal{M}}_{1,1}} \psi_1 = 1/24$ , so  $D_1 = 1/24$ . PENNER '85 computed  $V_{1,1}(0) = \zeta(2)$  implying  $D_0 = \pi^2/6$ . Else  $D_k = 0$  for  $k > 1$ . Four graphs of  $(0, 3)$ -type  $\Rightarrow A_{i_2, i_3}^{i_1} \neq 0$  iff  $i_* = 0$  ( $A_{0,0}^0 = 1$ )

- Q3: Is there a geometric object enumerated by tensor models?

5. TOWARDS BOREL SUMMABILITY IN  $1/N$ 

As a perspective for Borel summability in tensor models it is convenient to mention two essential techniques which together are the essence of the Loop Vertex Expansion V. RIVASSEAU *JHEP* '07. In L. FERDINAND, R. GURÄU, C.P & F. VIGNES-TOURNERET 2209.09045 BS in  $1/N$  is addressed for the cumulants of the vector model (defined below). The tools:

- The Hubbard-Stratonovich transformation

$$e^{-\frac{x^2}{2}} = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} dy e^{-\frac{y^2}{2} + ixy}$$

allows to transform the  $N$ -dimensional integral in the partition function of the  $O(N)$ -vector model

$$Z(g, 1/N, J) = \int_{\mathbb{R}^N} e^{-\frac{1}{2} \phi \cdot \phi - \frac{g}{8N} (\phi \cdot \phi)^2 - \sqrt{N} J \cdot \phi} \frac{d^N \phi}{(2\pi)^{N/2}}$$

into an integral where  $N$  appears only as a parameter, and not as the dimension of the integration domain, thus allowing for analytic continuation. With  $R(\sigma, z) = (1 - \sqrt{z\sigma})^{-1}$

$$Z(g, 1/N; J) = \int e^{\frac{N}{2} \log R(\sigma, g/N) + \frac{N}{2} R(\sigma, g/N) J \cdot J} \frac{e^{-\frac{\sigma^2}{2}} d\sigma}{\sqrt{2\pi}}$$

- Brydges-Kennedy-Abdesselam-Rivasseau proved

$$\Phi(\mathbf{1}) - \Phi(\mathbf{0}) = \sum_{\substack{\text{non-empty forests} \\ F \text{ with } n \text{ vertices}}} \int \frac{\partial \Phi}{\partial F} dF \quad (\text{BKAR})$$

$\mathbf{1}, \mathbf{0} \in \mathbb{R}^{\binom{n}{2}}$  the vectors with 1's and 0's as constant entries, respectively. The BKAR formula holds for any smooth function  $\Phi : \mathbb{R}^{\binom{n}{2}} \rightarrow \mathbb{C}$ , where the 'measure' over a forest and a 'forest' derivative are given by

$$dF := \prod_{e \in E(F)} du_e \quad (5a)$$

$$\frac{\partial \Phi}{\partial F} := \left\{ \left[ \prod_{(i,m) \in E(F)} \frac{\partial}{\partial x_{im}} \right] \Phi(x) \right\}_{x=w(F)} \quad (5b)$$

with parameters given by

$$w(F)_{ij} := \min\{u_e\}_{e \in E(F), e \text{ along the path } i \rightarrow j} \quad (5c)$$

and  $w(F)_{ij} = 0$  if such path does not exist. For instance, for  $n = 3$ , the forest expansion of  $\Phi(1, 1, 1)$  reads

$$\Phi(0, 0, 0) \quad \text{graph} \quad (6) \\ + \int_0^1 du_{12} \frac{\partial \Phi(x)}{\partial x_{12}} \Big|_{x=(u_{12}, 0, 0)} \quad \text{graph} \\ + \int_0^1 du_{23} \frac{\partial \Phi(x)}{\partial x_{23}} \Big|_{x=(0, u_{23}, 0)} \quad \text{graph} \\ + \int_0^1 du_{23} \frac{\partial \Phi(x)}{\partial x_{23}} \Big|_{x=(0, 0, u_{13})} \quad \text{graph} \\ + \int_0^1 \int_0^1 du_{23} du_{12} \frac{\partial^2 \Phi(x)}{\partial x_{12} \partial x_{23}} \Big|_{x=(u_{12}, u_{23}, \min\{u_{12}, u_{23}\})} \quad \text{graph} \\ + \int_0^1 \int_0^1 du_{23} du_{13} \frac{\partial^2 \Phi(x)}{\partial x_{23} \partial x_{13}} \Big|_{x=(\min\{u_{23}, u_{13}\}, u_{23}, u_{13})} \quad \text{graph} \\ + \int_0^1 \int_0^1 du_{13} du_{12} \frac{\partial^2 \Phi(x)}{\partial x_{12} \partial x_{13}} \Big|_{x=(u_{12}, \min\{u_{12}, u_{13}\}, u_{13})} \quad \text{graph}$$

- to have the cumulants, one needs the logarithm of this partition function. Performing Gaussian integration

$$Z(g, 1/N; J) = \sum_{n \geq 0} \frac{N^n}{(2)^n n!} \left[ \exp\left(\frac{\langle \partial, \partial \rangle_X(x)}{2N}\right) \prod_{i=1}^n \left\{ \log R(\sigma^{(i)}, g) + R(\sigma^{(i)}, g) J \cdot J \right\} \right]_{x_{ij}=1},$$

where  $[X(x)]_{ii} = 1$  and  $[X(x)]_{ij} = x_{ij}$  for  $i \neq j$ , one can use for the function in square-brackets BKAR-formula. Then  $\log Z$  will be the restriction from forests to trees.

- Q4: Are tensor models  $1/N$  Borel summable?

$$\begin{aligned}
 & \text{Surface with genus } g \text{ and boundary components } L = \{i_1, \dots, i_n\} \text{ label } \partial \\
 &= \sum_{m > 1} \text{Surface with genus } g \text{ and boundary components } i_1, \dots, i_m, i_{m+1}, \dots, i_n \\
 &+ \sum_{\text{stable}} \text{Surface with genus } g-1 \text{ and boundary components } i_1, i_2, \dots, i_n \\
 &+ \sum_{\substack{\text{stable} \\ h_1 + h_2 = g \\ J_1 \cup J_2 = L \setminus \{i_1\}}} \text{Surface with genus } g \text{ and boundary components } i_1, i_2, \dots, i_n
 \end{aligned}$$