Okinawa Seasonal Hazards

Okinawa Prefectural Office and Japan Meteorological Agency raise awareness on following common hazards

Prepared in March 2019

		1月	2月	3月	4月	5月	6月	7 月	8月	9月	10月	11月	12月		参考
		Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Note	Source (JPN)
ハブ	Habu														https://www.pref.okinawa.lg.jp/s ite/hoken/eiken/eisei/habutop.ht ml
ミツバチ	Honey Bees														https://www.pref.okinawa.lg.jp/s ite/hoken/eiken/eisei/hachi.html
アシナガバチ	Paper Wasp														https://www.pref.okinawa.lg.jp/s ite/hoken/eiken/eisei/hachi.html
スズメバチ	Hornet													スズメバチは年によっては冬でも巣の まわりに雌と雄が群がることがある。 Hornets may swarm around their nest during the winter season	https://www.pref.okinawa.lg.jp/s ite/hoken/eiken/eisei/hachi.html
タイワンキドク ガの幼虫	Orvasca taiwana larvae														https://www.pref.okinawa.jp/site /hoken/eiken/eisei/documents/28 page2.pdf
ハブクラゲ	Box Jellyfish*														https://www.pref.okinawa.jp/site /hoken/eiken/eisei/uminokikenns eibutu.html
台風	Typhoon**														http://www.jma.go.jp/jma/kisho u/nougyou/document/okinawa/o kinawa.pdf#page=33
梅雨	Rainy Season**														http://www.jma.go.jp/jma/kisho u/nougyou/document/okinawa/o kinawa.pdf#page=11
紫外線 (UV指標)	UV (UV Index)														https://www.jma- net.go.jp/okinawa/know/memo/p df/sougou/201806bosai.pdf
熱中症	Heatstroke													Measures from first week of June till last week of September (19 weeks)	https://www.pref.okinawa.jp/site /hoken/chiikihoken/shippei/heatf lyer/heatstroke/documents/19- l.pdf
強風・高波	Strong wind High wave													北風 Strong northern winds	http://www.jma- net.go.jp/okinawa/know/kaiyo/te nko.html



^{*}Most reported cause of marine animal injury. There are other marine animals that can cause injuries as well.

^{**}Some lead to secondary hazards. Example: excessive rain may also cause landslides